

Anti-pStat3 [Y705]-158Gd

Catalog: 3158005A

Package size: 50 tests

Storage: Store product at 4 °C. Do not freeze.

Cross-reactivity: Mouse, Human

Clone: 4/P-STAT3


Isotype: Mouse IgG2a

Formulation: Antibody stabilizer with 0.05% sodium azide

Technical Information

Validation: Each lot of conjugated antibody is quality control-tested by CyTOF[®] analysis of stained cells using the appropriate positive and negative cell staining and/or activation controls.

Recommended usage: The suggested use is 1 µL for up to 3 × 10⁶ live cells in 100 µL. It is recommended that the antibody be titrated for optimal performance for each of the desired applications.



pStat3 [Y705] (4/P-STAT3)-158Gd

Human Jurkat T cells were incubated for 15 minutes in media alone (bottom) or with pervanadate (top). Cells were then fixed, permeabilized and stained with 158Gd-anti-pStat3 [Y705] (4/P-STAT3).

Description

Members of the STAT (signal transducers and activators of transcription) family are important intracellular messengers of cytokines and growth factor signaling. Seven mammalian STATs have been identified: STAT1-4, 5a, 5b, and 6. STAT proteins are activated by tyrosine phosphorylation, which causes dimerization and translocation to the nucleus, where the STAT dimer acts as a transcription factor. JAK-mediated phosphorylation of Tyr705 on STAT3 occurs in response to many cytokines and growth factors including interferon-alpha, EGF, IL-5, IL-6, G-CSF and HGF. Activated STAT3 promotes transcription of genes that mediate cell growth and differentiation.

References

Bandura, D. R., et al. Mass Cytometry: Technique for Real Time Single Cell Multitarget Immunoassay Based on Inductively Coupled Plasma Time-of-Flight Mass Spectrometry. *Analytical Chemistry* 81 (2009): 6,813–22.

Ornatsky, O. I., et al. Highly Multiparametric Analysis by Mass Cytometry. *Journal of Immunological Methods* 361 (2010): 1–20.

Bendall, S. C., et al. Single-Cell Mass Cytometry of Differential Immune and Drug Responses Across a Human Hematopoietic Continuum. *Science* (2011): 687–96.

For technical support visit www.fluidigm.com/support.

North America +1 650 266 6100 | Toll-free: +1 866 358 4354 (US/CAN) | support.northamerica@fluidigm.com **Europe** +44 1223 859941 | support.europe@fluidigm.com

China (excluding Hong Kong) +86 21 3255 8368 | techsupportchina@fluidigm.com **Japan** +81 3 3662 2150 | techsupportjapan@fluidigm.com

All other Asian countries +1 650 266 6100 | techsupportasia@fluidigm.com **Central and South America** +1 650 266 6100 | techsupportlatam@fluidigm.com

For Research Use Only. Not for use in diagnostic procedures.

Information in this publication is subject to change without notice. **Safety data sheet information:** www.fluidigm.com/sds. **Patent and license information:** www.fluidigm.com/legalnotices. **EU's WEEE directive information:** www.fluidigm.com/compliance. Fluidigm, the Fluidigm logo, and CyTOF are trademarks or registered trademarks of Fluidigm Corporation in the United States and/or other countries. © 2017 Fluidigm Corporation. All rights reserved. 02/2017